


Nutrient Deficiency **ID Guide** *for Hemp*



Published by the Ministry of Agriculture, Food and Rural Affairs

© King's Printer for Ontario, 2023

Toronto, ON

ISBN 978-1-4868-7132-2 (Print)

ISBN 978-1-4868-7133-9 (PDF)

Cette publication est aussi disponible en français.

Nutrient Deficiency **ID Guide** *for Hemp*



Nutrient Deficiency ID Guide for Hemp

Editor

Jim Todd, Ontario Ministry of Agriculture, Food and Rural Affairs (OMAFRA).

Graphic Design/Art Direction

Andrea Vieira (OMAFRA)

Acknowledgments

The author would like to thank Rachel Riddle, University of Guelph, for technical assistance, A & L Laboratories for in-kind contributions towards nutrient analysis and the Knowledge Translation and Transfer (KTT) Funding Program, Ontario Agri-Food Innovation Alliance for funding this project (grant # UG-KTTM-2020-100738).

*If you require technical or business information,
contact the Agricultural Information Contact Centre
at 1-877-424-1300 or ag.info.omafra@ontario.ca*

Contents

| | |
|-----------------------------------|----|
| Introduction | 3 |
| Nitrogen (N) | 6 |
| Phosphorus (P) | 8 |
| Potassium (K) | 10 |
| Calcium (Ca) | 12 |
| Magnesium (Mg) | 14 |
| Sulphur (S) | 16 |
| Boron (B) | 18 |
| Chlorine (Cl ⁻) | 20 |
| Copper (Cu) | 22 |
| Iron (Fe) | 24 |
| Manganese (Mn) | 26 |
| Molybdenum (Mo) | 28 |
| Zinc (Zn) | 30 |
| Glossary | 32 |
| Citations | 34 |

Table 1. Form and mobility of nutrients and micronutrients in soil and plants.

| Nutrients | Mobility in Soil | Plant available forms in soil | Mobility in plant |
|----------------------------|------------------|---|-------------------|
| Primary Nutrients | | | |
| Nitrogen | medium | ammonium ion (NH_4^+) | high |
| Nitrogen | high | nitrate ion (NO_3^-) | high |
| Phosphorus | low | phosphate ion (H_2PO_4^- , HPO_4^{2-}) | high |
| Potassium | low–medium | potassium ion (K^+) | high |
| Secondary Nutrients | | | |
| Calcium | low | calcium ion (Ca^{2+}) | low |
| Magnesium | low | magnesium ion (Mg^{2+}) | high |
| Sulphur | medium | sulphate ion (SO_4^{2-}) | low–medium |
| Micronutrients | | | |
| Boron | high | boric acid ($\text{B}(\text{OH})_3^0$, borate ion (H_2BO_3^-) | low–medium |
| Chlorine | high | chloride ion (Cl^-) | high |
| Copper | low | cupric ion (Cu_2^+) | low |
| Iron | low | ferrous ion (Fe^{2+}), ferric ion (Fe^{3+}) | low |
| Manganese | low | manganous ion (Mn^{2+}) | low |
| Molybdenum | low–medium | molybdate ion (MoO_4^{2-}) | medium–high |
| Zinc | low | zinc ion (Zn^{2+}), zinc hydroxide $\text{Zn}(\text{OH})_2^0$ | low |

Source: OMAFRA, Publication 611, *Soil Fertility Handbook*, Table 3-3.



Introduction

Increases in hemp and cannabis production in Ontario have resulted in numerous inquiries from growers on how foliar symptoms relate to various nutrient deficiencies. Resources for visual identification of nutrient deficiency symptoms in hemp are limited. Those currently available typically have old, poor-quality photos that do not cover the full nutrient spectrum or the progressive development of deficiency symptoms seen in the field.

The ability to identify nutrient deficiencies quickly and accurately allows growers to improve nutrient use efficiency, crop health and overall yields of marketable product. This pictorial guide was developed for growers to better identify foliar symptoms caused by nutrient deficiencies in *Cannabis sativa* plants.

Female industrial hemp plants (cv. Finola) were grown under standard greenhouse conditions. Supplemental lighting was provided to ensure plants did not convert to reproductive mode. Individual solutions, each lacking one of the 13 macronutrient or micronutrients (Table 1), and one complete nutrient treatment based on Hoagland's solution corrected to pH 6.5 were applied to plants for the duration of the project. A pH of 6.5 was chosen because all nutrients are available at that pH (Figure 1). Photos were taken of deficiency symptoms at multiple stages for each nutrient and a companion illustration of the types of symptoms that occur with nutrient deficiencies is provided (Figure 2).




Figure 1. Effect of pH on the root availability of the essential elements in soil. Blue denotes the ideal soil pH for the majority of plants (slightly acidic).
 Source: Wikimedia Commons^[1]




Figure 2. Illustration of the typical symptoms that occur in a leaf when a plant experiences nutrient deficiencies.

Nitrogen (N)

Primary Nutrient

Role

Nitrogen is involved in many plant processes and structures. Compared to other nutrients, it is required in large amounts. Nitrogen is a main component of amino acids, which form proteins within the plant. Enzyme proteins are important in several plant processes, particularly those which impact growth and yield. Protein is usually highest in the harvested part of the plant hence it often is an important item in the nutritional value of the crop. Nitrogen has an important role in the production of chlorophyll, which gives the green colour in plants. Chlorophyll is responsible for the conversion of sunlight to energy needed by the plant through the process of photosynthesis.

Mobility

Nitrogen is a mobile element in the plant.

Impact of pH

Availability decreases under highly acidic or alkaline conditions.

Deficiency Symptoms

Plants suffering nitrogen deficiency are usually a pale green to yellow, with symptoms first occurring in the older leaves which may show some red or purple tinting, particularly in the veins and petioles of broadleaf plants. Plants may show reduced vigour.



Figure 3. Progression of leaf yellowing due to nitrogen deficiency in young to old leaves (left to right).



Figure 4. Plant showing older mature leaves in the lower half of the plant that are yellow due to nitrogen deficiency.

Phosphorus (P)

Primary Nutrient

Role

Like nitrogen, phosphorus is an important factor in many plant metabolic processes such as photosynthesis and respiration, energy storage and transfer (ATP), protein and carbohydrate metabolism, and cell division and enlargement. It is also integral to the structure of DNA and is an important component of cell membranes. The effect of the availability and supply of phosphorus on these plant processes and structures is reflected in specific aspects of crop growth. When phosphorus is limiting, plants adjust by shifting more resources to root production and less to top growth.

Mobility

Phosphorus is a mobile element in the plant.

Impact of pH

Availability begins to decrease below pH 6.

Deficiency Symptoms

Purpling in plants is associated with phosphorus deficiency, but this symptom is unreliable. The production of anthocyanin, which creates the purple colour, is a standard stress response. Many other factors can induce purpling. Common phosphorus deficiency symptoms include slow-growing plants with dark-green leaves in early deficiency stages of older leaves, delayed plant maturity, and spindly, small leaflets. Older leaves typically progress from dark green to bronze tinting with interveinal and marginal necrosis (browning). There may also be a decrease in the number and size of buds or flowers.



Figure 5. Progression of phosphorus deficiency symptoms from youngest (left) to oldest (right) leaves.



Figure 7. Dark green leaves are a common phosphorus deficiency symptom.



Figure 6. Extreme phosphorus deficiency will result in stunted plants showing reduced vigour.



Figure 8. Purpling of plant stems and petioles induced by phosphorus deficiency.

Potassium (K)

Primary Nutrient

Role

Potassium promotes formation of structural components including lignin and cellulose, which play a major role in stalk strength and lodging resistance. It influences the uptake of carbon dioxide, photosynthesis and the regulation of stomatal opening in the leaves, water uptake by roots, starch and sugar content and cell integrity, and aids in disease and insect resistance.

Mobility

Potassium is a mobile element in the plant.

Impact of pH

Availability decreases below pH 6 and is reduced between pH 7.6 to 8.4.

Deficiency Symptoms

Since potassium is mobile within the plant, deficiency symptoms usually appear first on older leaves, often as a chlorosis (yellowing) or necrosis of the leaf margins known as 'marginal scorch'. Older leaves often curl downwards. Potassium deficiency may also cause shorter internodes.



Figure 9. Development of chlorosis and necrosis of the leaf margins from young (left) to older leaves (right) because of potassium deficiency.



Figure 10. Leaf clusters showing symptoms of chlorosis or necrosis of the leaf margins due to potassium deficiency.

Calcium (Ca)

Secondary Nutrient

Role

Calcium is important in the stabilization of the cell wall and is involved in the metabolism and formation of the cell nucleus.

Mobility

Calcium is an immobile element in the plant. Uptake is depressed by ammonium based nitrogen, excessive potassium, magnesium, manganese and aluminum.

Impact of pH

Availability declines below pH 6.5.

Deficiency Symptoms

Plants demonstrate slow growth and reduced size eventually resulting in no further elongation of the growing points and finally death of the growing points. Leaves typically turn pale green during early onset of calcium deficiency. Leaves, particularly new growth, progress with patches of yellow to brown interveinal spotting and necrosis. Leaf curling can also occur.



Figure 11. Young leaves will develop yellow-brown (necrotic) spotting and curled leaf margins as a result of calcium deficiency.



Figure 12. Leaves can take on unusual shapes with prolonged calcium deficiency.



Figure 13. Yellow-brown necrotic spots will become extensive with prolonged calcium deficiency.

Magnesium (Mg)

Secondary Nutrient

Role

Magnesium is required for the synthesis of chlorophyll for photosynthesis. It also aids phosphate movement and is a component of enzymes involved in plant respiration and DNA/RNA synthesis.

Mobility

Magnesium readily moves through the plant.

Impact of pH

Optimally available between pH 6.5 and 8.5.

Deficiency Symptoms

Magnesium is a mobile element, so older leaves are the first to display signs of deficiency. These typically include interveinal chlorosis or reddening followed by marginal leaf necrosis, leaf curling and development of necrotic patches. Magnesium deficiency symptoms may appear similar to potassium and manganese deficiency symptoms however, manganese deficiency symptoms usually occur first in younger leaves compared to magnesium and phosphorus deficiency symptoms first occurring in older leaves.



Figure 14. Early development of magnesium deficiency symptoms. First signs of injury appear on the older leaves. Interveinal tissues begin to exhibit chlorosis. Magnesium deficiency may have been occurring for several weeks before symptoms are visible.



Figure 15. Mid-development of magnesium deficiency symptoms: Cell breakdown in chlorotic tissues advances to tissue necrosis resulting in patches of brown, dead tissue.



Figure 16. Late development of magnesium deficiency symptoms: Progression from early chlorosis to cell death as leaves age is shown. Leaf curling is a common feature of magnesium deficiency. Damage is primarily located in the older leaves at the base of the plant.



Figure 17. Late development of magnesium deficiency symptoms showing lower leaves that are necrotic and curled.

Sulphur (S)

Secondary Nutrient

Role

Sulphur is a constituent of two of the 21 amino acids that form proteins. It also helps develop enzymes and vitamins, is involved in nitrogen fixation in legumes, aids in seed production and is needed for chlorophyll formation.

Mobility

Sulphur is an immobile element in the plant.

Impact of pH

Availability decreases below pH 6.0.

Deficiency Symptoms

Plants suffering sulphur deficiency are usually a pale green and may show some red or purple tinting similar to nitrogen deficiency symptoms. The younger leaves may be more chlorotic than the older leaves due to the sulphur being immobile in the plant. Stems can become woody in appearance with shortened internodes.



Figure 18. Young leaves showing chlorosis in response to sulphur deficiency.



Figure 19. Increased stem woodiness may also occur because of sulphur deficiency.



Figure 20. Plants showing chlorosis in upper leaves (left panel) and general paleness in new growth (right panel) due to sulphur deficiency.



Figure 21. Progression of chlorosis from moderate (left) to severe (right) because of sulphur deficiency.

Boron (B)

Micronutrient

Role

Boron plays an important role in the structural integrity of cell walls, fruit set, seed development and carbohydrate and protein metabolism.

Mobility

Boron is an immobile element in the plant.

Impact of pH

Availability declines above pH 7.5.

Deficiency Symptoms

Symptoms first appear on younger tissues, commonly as distorted (curled) leaves that become thicker with age. Interveinal chlorosis followed by necrosis is evident. In general, vegetative growth will be poor. Growing points typically die back, and buds or flowers may be deformed or die.



Figure 22. Symptoms of boron deficiency appear in the younger leaves and include interveinal chlorosis, distortion, and thickening.



Figure 23. Distortion of the younger leaves is a common symptom of boron deficiency.

Chlorine (Cl⁻)

Micronutrient

Role

Chlorine in plants plays an important role in stomatal regulation and water flow. Chlorine is also involved in photosynthesis.

Mobility

Chlorine is readily soluble and highly mobile within the plant.

Impact of pH

Mobility not affected by pH.

Deficiency Symptoms

Plants will demonstrate reduced growth. Wilting or limpness of growing points and young leaves is typical of early chlorine deficiency symptoms followed by the development of chlorotic and necrotic spots. Younger leaves may also demonstrate leaf cupping. Bronzing of the leaves occurs after prolonged deficiency.



Figure 24. Progression from yellow chlorotic spots (upper left panel) to brown necrotic patches (right and lower panels) as a result of chlorine deficiency.



Figure 25. Bronzing of the leaf surface will occur with prolonged chlorine deficiency.

Copper (Cu)

Micronutrient

Role

Copper plays a role in chlorophyll production and proper enzyme function.

Mobility

Copper is an immobile element in the plant.

Impact of pH

Availability declines above pH 8.0.

Deficiency Symptoms

Leaves, particularly new growth, will develop a dark blueish-green colour and show signs of wilting. With time, leaves become chlorotic and may begin to curl.



Figure 26. Progression of chlorosis to necrosis (left to right) in young to old leaves due to copper deficiency.



Figure 27. Blueish-green leaves that are beginning to curl are symptomatic of early stages of copper deficiency.



Figure 28. With time, leaves begin to droop in response to a lack of copper.

Iron (Fe)

Micronutrient

Role

Iron has a number of functions within plants. It is a catalyst in the formation of chlorophyll and is required for plant respiration and functions in the formation of some proteins.

Mobility

Iron is an immobile element in the plant.

Impact of pH

Availability declines above pH 7.5.

Deficiency Symptoms

The first signs of iron injury appear on the younger leaves which show interveinal chlorosis or striping along the entire length of the leaf. Interveinal chlorosis progressively develops from light green to light yellow or white, eventually showing the appearance of brown patches of dead tissue. Leaves stay alive during iron deficiency and only die after prolonged iron deprivation. In severe cases, plant growth will be stunted.



Figure 29. A lack of iron results in young leaves showing interveinal chlorosis starting from the base of the leaf.



Figure 30. Development of necrotic spots as chlorosis progresses in response to iron deficiency.



Figure 31. Iron deficiency symptoms occur on leaves in the upper half of the plant.



Figure 32. Prolonged iron deficiency results in severe necrosis leading to leaf death.

Manganese (Mn)

Micronutrient

Role

Many enzyme systems require manganese for proper function, including those involved in the distribution of plant growth regulators within the plant. It is involved in photosynthesis and chlorophyll production.

Mobility

Manganese is an immobile element in the plant.

Impact of pH

Availability declines above pH 7.5.

Deficiency Symptoms

The first signs of injury appear on the younger leaves. Leaf margins and veins may look darker green than the rest of the leaf. Interveinal yellowing progressively develops into the appearance of necrotic speckling or patches of dead tissue. Manganese deficiency symptoms may appear similar to potassium and magnesium deficiency symptoms however, manganese deficiency symptoms usually occur first in younger leaves compared to magnesium and phosphorus deficiency symptoms first occurring in older leaves.



Figure 33. Progressive development of manganese deficiency symptoms. The youngest leaf in the upper left shows darker green leaf margins. Interveinal yellowing develops as the leaf ages (upper right and lower panels).



Figure 34. Progressive development of manganese deficiency symptoms from youngest leaf on the left to oldest leaf on the right.

Molybdenum (Mo)

Micronutrient

Role

Molybdenum plays an important role in nitrogen metabolism within the plant as it is an essential component of nitrate reductase, the enzyme that converts nitrate (NH_3^-) to ammonium (NH_4^+). It also plays a role in pollen viability and seed production.

Mobility

Molybdenum is a mobile element in the plant.

Impact of pH

Availability increases above pH 7.0.

Deficiency Symptoms

A lack of molybdenum impacts nitrogen metabolism, so molybdenum deficiency symptoms may be similar to nitrogen deficiency symptoms including yellowing, plant stunting, scorching of the leaf margins and cupping or rolling of leaves. Curling or deformity of younger leaves is commonly referred to as 'whiptail' in many plant species. Deficiency symptoms first occur in older leaves and then progress to the younger, expanding leaves. Both younger and older leaves may develop chlorotic spots that eventually perforate but leave the rest of the leaf tissue green for a prolonged time period.



Figure 35. Development of yellowing from young leaf (top) to older leaves (middle and bottom) because of molybdenum deficiency.



Figure 36. Leaf displaying cupping due to molybdenum deficiency.



Figure 37. Leaves displaying cupping and rolling due to molybdenum deficiency.

Zinc (Zn)

Micronutrient

Role

Zinc is important in early plant growth and in grain and seed formation. It plays a role in chlorophyll and carbohydrate production.

Mobility

Zinc is an immobile element in the plant.

Impact of pH

Availability declines above pH 8.0.

Deficiency Symptoms

Interveneal chlorosis is evident in the youngest tissues which may appear as pale to white bands between the leaf margin and mid-vein in the lower part of the leaf. In severe cases, newly emerged leaves can be completely white. In addition, the developing leaves may be thin with distorted margins.



Figure 38. Yellow to white interveinal chlorosis in the younger leaves is a common symptom of zinc deficiency.



Figure 39. Interveinal chlorosis in the younger leaves is a common symptom of zinc deficiency.



Figure 41. Zinc is an immobile nutrient, so the initial symptoms occur at the top the plant.



Figure 40. Thin, strappy younger leaves may also result from zinc deficiency.



Figure 42. Interveinal chlorosis initially occurs in the lower half of the leaf when zinc is deficient.

Glossary

Bronzing: A reddish-brown discoloration or flecking of plant tissues, especially leaves, in response to damage from a pest (e.g. mite feeding) or environmental stress (e.g. sunscald, excess ozone, nutrient deficiency). Leaves may also appear curled and pale green in colour.

Chlorosis: Bleaching (yellowing) of plant tissues due to a loss of chlorophyll.

Intraveinal: Occurring within the leaf veins.

Internode: The part of a stem or rhizome between any two nodes.

Interveinal: Area located between the veins that occur in leaves

Margin (leaf): Area along the edge of the leaf.

Micronutrient: Micronutrients are as important as the primary and secondary nutrients but are needed in much smaller quantities by the plant and are often less prevalent in the soil.

Mobile Nutrient: Any nutrient that can move freely within the plant. Mobile nutrients move to the areas of new growth, which can result in older tissues displaying deficiency symptoms.

Necrosis: Premature tissue death resulting from disease, injury or physiological stress (e.g., nutrient deficiency, ozone etc.). Necrosis can occur within the body of the leaf, along the leaf margins, or in both locations.

Necrotic Speckling/Stippling: Numerous small, white or bronzed puncture marks on the leaf.

Petiole: The stalk of a leaf, attaching the blade to the stem.

Primary Nutrient: A primary nutrient, or macronutrient, is required by plants in large quantities for basic plant growth and development. The six nutrients that fall into this category are carbon, hydrogen, oxygen, nitrogen, phosphorus and potassium. Plants acquire carbon, hydrogen and oxygen from the air and water. The remaining macronutrients must be obtained from the soil. Fertilizer, manure, nitrogen fixation and mineral weathering replenish soil nutrients. Primary nutrients most frequently limit plant growth.

Secondary Nutrient: Calcium, magnesium and sulphur are required in moderate amounts. They are usually classified as secondary nutrients because they are less likely to limit crop growth. These nutrients are usually present in the soil in adequate amounts, although fertilization may be required for certain crops.

Scorch (leaf): Sudden leaf death or browning, either interveinally or at the margin that occurs when plants have difficulty taking up water or have sudden exposure to full sunlight following a wet, cloudy period.

Stunting: Reduced plant growth due to a pest or a lack of water, nutrients or other necessity for plant development.

Tipburn: Yellowing and/or necrosis of the leaf tissue at the apex.

Citations

- 1 Wikimedia Commons, *File:Soil pH effect on nutrient availability.svg*, https://commons.wikimedia.org/w/index.php?title=File:Soil_pH_effect_on_nutrient_availability.svg&oldid=634347869#filelinks. Accessed March 14, 2023, [Creative Commons Attribution 4.0 International license](#).

References

Ontario Ministry of Agriculture, Food and Rural Affairs, Publication 611, *Soil Fertility Handbook*, Queen's Printer for Ontario, 2019.

Hewitt, E.J. and T.A. Smith. (1975). *Plant mineral nutrition*. The English Universities Press Ltd. London, U.K. 1975. Hoagland, D.R. and D.I. Arnon. 1950. *The water culture method for growing plants without soil*. Circ. 347. California Agr. Expt. Stn., Univ. of California, Berkeley.

Bould, C., Hewitt, E.J., and Needham, P. *Diagnosis of Mineral Disorders in Plants*, Volume 1: Principles. Her Majesty's Stationary Office, London, U.K. 1983.

Photo Credits

- | | |
|------|---|
| FC | Hemp leaf at sunset, Shutterstock, 207279334 |
| p. 3 | Hemp field panorama, Shutterstock, 1786911731 |
| BC | Hemp leaf at sunset, Shutterstock, 207279334 |

This guide is published for informational purposes only. The Province of Ontario, as represented by the Ontario Ministry of Agriculture, Food and Rural Affairs (OMAFRA), disclaims any express or implied warranties related to the use of this guide, including all contents, any link to or the contents of any third-party site or source, including, without limitation, warranties of non-infringement or of fitness for any particular purpose.

In no event shall the Province of Ontario or its directors, officers, employees, servants or agents accept any liability for any failure to keep the guide's contents up to date or for any errors or omissions within it or in any link or third party site or sources that may be referenced within it or for any damages (including, without limitation, damages for loss of profits, business interruption, loss of information, or direct, indirect, incidental, special consequential or punitive damages), whatsoever arising out of or related to the use of or inability to use this guide (including all contents), any link or any third party site or works, whether under contract, in tort or under any other basis of liability. It is the user's responsibility to ensure they have chosen the best course of action for their own particular circumstances.

The contents of this guide (including, without limitation, the graphics, icons, and its overall appearance) are the property of the Province of Ontario. The Province does not waive any of its proprietary rights therein including, but not limited to, copyrights, trademarks and other intellectual property rights.

No user of this guide may sell, republish, print, download, copy, reproduce, modify, upload, post, transmit or distribute in any way any portion of this guide or its contents without the prior written consent of the Province, except for reasonable printing, downloading and copying for the private information and use of the user. The availability of any of the contents of this guide shall under no circumstance constitute a transfer or waiver of any copyrights, trademarks or other intellectual property rights of the Province to any user or to any third party.

